

nature genetics

JSM
96-127
SIBL

Nature genetics.
v. 25
no. 4
Aug 2000

U.S. 20

SIBL

JD

volume 25 no. 4

august 2000


101633
AUTO SCIENCE 3-DIGIT 101
101633 SCIENCE 005 DEC00 02000
NY PUBLIC LIBRARY
NY PUBLIC LIBRARY
SCIENCE LIBRARY
SCIENCE LIBRARY
GRAND CENTRAL STATION
PO BOX 5123
NEW YORK NY 10163-2233
0000004/0037

Poln takes on
transversions


Selection on
BRCA1

Mouse
mutagenesis

Gene for Robinow
syndrome



<http://genetics.nature.com>



Cover art by
Mitsuko Hoshino

nature genetics

volume 25 no. 4

august 2000

editorial

- The end of the beginning 363

news & views

- Hunting in the calm before the storm 365
Gillian Bates & James Eberwine **SEE ALSO** 385

- Tailoring vectors through DNA shuffling 366
David T Curiel **SEE ALSO** 436

- Subdividing the metamer: one signal, two outcomes 368
Claudio D Stern **SEE ALSO** 390

- TOUCHINGbase 371

correspondence

- Of sex and gender 373
D Haig

- Genetic homogeneity of Icelanders: fact or fiction? 373
E Arnason, H Sigurgislason & E Benedikz

brief communications

- Hot-stop PCR: a simple and general assay for linear quantitation of allele ratios 375
H Uejima, M P Lee, H Cui & A P Feinberg

- The *NOTCH4* locus is associated with susceptibility to schizophrenia 376
J Wei & G P Hemmings

Nature Genetics (ISSN 1061-4036) is published monthly by Nature America Inc., headquartered at 345 Park Avenue South, New York, NY 10010-1707. Editorial Office: 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726 9314, Fax: (212) 545 8341. North American Advertising: 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726 9200, Fax: (212) 696 9006. European Advertising: *Nature Genetics*, Porters South, Crinan Street, London N1 9XW, UK. Tel: 44 171 833 4000, Fax: 44 171 843 4996. New subscriptions, renewals, changes of address, back issues and all customer service questions in the Americas should be addressed to: *Nature Genetics* Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Tel: (800) 524 0384, Direct Dial (615) 377 3322, Fax: (615) 377 0525. Outside the Americas: *Nature Genetics*, Macmillan Magazines Ltd., Houndsmill, Brunel Road, Basingstoke, RG21 6XS, U.K. Tel: +44-(0)1256-329242, Fax: +44-(0)1256 812358. Email: subscriptions@nature.com. Annual subscription rates: North America: US\$650 (institutional/corporate), US\$199 (individual making personal payment). Canada add 7% for GST, BN: 14091 1595 RT; U.K./Europe: £475 (institutional/corporate), £110 (individual making personal payment), £99 (student); Rest of world (excluding Japan): £520 (institutional/corporate), £235 (individual making personal payment) £110 (student); Japan: Contact Nature Japan K.K., Shin-Mitsuke Building 3F, 3-6 Ichigaya Tamachi, Shinjuku-ku, Tokyo 162-0843. Periodicals postage paid at New York, NY 10010, and at additional mailing offices. POSTMASTER: Send address changes to *Nature Genetics* Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Executive Officers of Nature America Inc.: Nicholas Byam Shaw, Chairman of the Board; Jan Velterop, President; Edward Vallis, Secretary-Treasurer. Published in Japan by Nature Japan K.K., Shin-Mitsuke Building 3F, 3-6 Ichigaya Tamachi, Shinjuku-ku, Tokyo 162-0843, Japan. Printed by Cadmus Journal Services, Richmond, VA, US. Copyright ©2000 Nature America Inc.

brief communications




Fig. 2 Comparison of regular PCR and hot-stop PCR for quantitation of allele ratios. **a**, Representative result at 35 cycles of PCR, comparing regular PCR with hot-stop PCR at varying ratios of *IGF2* template, with primers P3 and P4 as described³. A allele, 294 bp, undigested by *Apa*I; B allele, 229 bp, digested by *Apa*I; DC, internal digestion control for this experiment (179 bp undigested, 126 bp digested). **b**, Linearity and accuracy of hot-stop PCR compared with regular PCR. Three independent experiments were done and the averages of allele ratios from regular PCR and hot-stop PCR are indicated by green circles and blue triangles, respectively, and ratios from regular PCR are indicated by red squares. Red squares represent the theoretical values. **c**, Representative results of analysis of *IGF2* imprinting by hot-stop PCR of normal and tumour samples. Normal tissue was from a fetal kidney that showed complete imprinting of *IGF2*. Tumours showed partial or complete LOI. LOI in the cancer represented by the fourth panel is evident despite a 2:1 genomic DNA allele ratio due to aneuploidy. G, + and - indicate that the PCR templates were from genomic DNA, cDNA (RT+) or cDNA (RT-), respectively.

PCR (Fig. 2a,b). In contrast, using hot-stop PCR, the A:B ratio precisely reflected the theoretical ratio (Fig. 2a,b). As expected, the allele ratio measured by hot-stop PCR was independent of PCR cycle number, amount of template or PCR amplification efficiency, whereas regular PCR was sensitive to these variables.

We also analysed the imprinting status of *IGF2* in normal and tumour cells. Normal imprinting, LOI and partial LOI were observed and quantified. The assay was also sensitive enough to quantify genomic allele ratio in aneuploid cells, allowing correction of allele-specific expression ratio for the allele-specific genomic template ratio (Fig. 2c). These results do not contradict previous studies examining ratios of allele-specific expression, provided that those experiments were performed with careful attention to limiting cycle number and avoiding high template concentration, but

they do illustrate the possibility of errors introduced by regular PCR. We expect that hot-stop PCR will allow for more consistent and reliable measurement of small differences in these ratios, an important issue for genes that show subtle tissue-specific changes in genomic imprinting¹⁻³.

Acknowledgements

We thank P. Onyango and J. Ravenel for discussions. This work was supported by grants from the National Institutes of Health to A.P.F.

Hiroshi Uejima^{1,2*}, Maxwell P. Lee^{1,2*}, Hengmi Cui^{1,2} & Andrew P. Feinberg^{1,2,3}

*These authors contributed equally to this work.

¹Institute of Genetic Medicine, and Departments of ²Medicine, ³Molecular Biology & Genetics and Oncology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA. Correspondence should be addressed to A.P.F. (e-mail: afeinberg@jh.edu).

The *NOTCH4* locus is associated with susceptibility to schizophrenia

Linkage disequilibrium mapping of the MHC region in 80 British parent-offspring trios showed that *NOTCH4* was highly associated with schizophrenia. The A→G substitution in the promoter region and the (CTG)_n repeat in exon 1 of

NOTCH4 may be candidate sites conferring susceptibility to schizophrenia.

Linkage studies indicate that chromosome 6p may contain a susceptibility locus for schizophrenia¹⁻⁴, although the evidence is not strong enough to draw firm conclusions. It may be that a gene of moderate or small effect rather than one of major effect is involved. Because the human major histocompatibility complex (MHC) region is mapped to 6p21.3 and its complete genomic DNA sequence is now available, we conducted the transmission disequilibrium test⁵ (TDT) with densely spaced DNA markers to search for a schizophrenia susceptibility gene within the MHC region.

We examined 80 parent-offspring trios of British descent, consisting of fathers,

1. Lee, M.P., Hu, R.-J., Johnson, L.A. & Feinberg, A.P. *Nature Genet.* **15**, 181-185 (1997).

2. Qian, N. et al. *Hum. Mol. Genet.* **6**, 2021-2029 (1997).

3. Cui, H., Haron, I.L., Ohlsson, R., Hamilton, S.R. & Feinberg, A.P. *Nature Med.* **4**, 1276-1280 (1998).

4. Anonymous. *ABI Prism DNA Sequencing Chemistry Guide* (Perkin-Elmer, Foster City, 1995).

5. Mansfield, E.S. *Hum. Mol. Genet.* **2**, 43-50 (1993).

6. Ohlsson, R. et al. *Genomic Imprinting: Causes and Consequences* (Cambridge: University Press, Cambridge, 1995).

7. Mashal, R.D., Koontz, J. & Sklar, J. *Nature Genet.* **9**, 177-183 (1995).

8. Takeda, S., Ichii, S. & Nakamura, Y. *Hum. Mutat.* **2**, 112-117 (1993).

9. Pushnova, E.A. & Zhu, Y.S. *Anal. Biochem.* **260**, 24-29 (1998).

10. Yun, K., Soejima, H., Merric, A.E., McCall, J.L. & Reeve, A.E. *J. Pathol.* **187**, 518-522 (1999).

11. Greenwood, A.D. & Burke, D.T. *Genome Res.* **6**, 336-348 (1996).

12. Rainier, S. et al. *Nature* **362**, 747-749 (1993).

13. Ogawa, O. et al. *Nature* **362**, 749-751 (1993).

brief communications

Table 1 • LD analysis of the class II and class III regions of the MHC in schizophrenia

Locus	Accession	Marker	Heterozygosity	P value ^a
AC004180	AC004180	(GT)n	0.85	0.145
HSY14768	Y14768	(CA)n	0.82	0.16
DJ201G24	AF129756	(GT)n	0.82	0.311
HSA012008	AJ012008	(CA)n	0.67	0.463
HSMHC3W36A	U89337	(TTG)n	0.73	0.5
HSMHC3A5	U89335	(TAA)n ^b	0.74	0.00017
		SNP1 ^c	0.16	0.267
		SNP2 ^d	0.26	0.002
		(CTG)n ^e	0.79	0.000036
		(TTAT)n ^f	0.51	0.393
HS107715	AL034394	(CT)n	0.66	0.356
HS172K2	Z84814	(GT)n	0.75	0.5
HSDV19	Z84490	(CA)n	0.73	0.372
HSEVMHC	X87344	(TAA)n	0.59	0.016
HSO14	Z84497	(GT)n	0.76	0.5
HS1033B10	AL031228	(CCT)n	0.53	0.5
HSICK721Q	AL021366	(GT)n	0.83	0.108

^aThe TDTLKE program for the McNemar test was applied to analyse genotyping data¹¹. The significance level of a P value was set at 0.003 as 17 markers were tested. ^bFrequencies of 8 transmitted alleles were 1, 53, 22, 22, 5, 7, 5 and 4, and of 8 non-transmitted alleles, 1, 21, 35, 32, 7, 13, 4 and 6. ^cThe T→C substitution creates a *MspI* site. Frequencies of the 2 transmitted alleles were 30 and 22. ^dThe A→G substitution also creates a *MspI* site. Frequencies of the 2 transmitted alleles were 27 and 55. ^eFrequencies of 7 transmitted alleles were 1, 13, 23, 63, 8, 17 and 1, and of 7 non-transmitted alleles, 1, 18, 47, 23, 12, 22 and 3. ^fFrequencies of 6 transmitted alleles were 5, 0, 31, 20, 11 and 2, and of 6 non-transmitted alleles, 9, 3, 24, 23, 9 and 1. Only 68 of 80 parent-offspring trios were available for analysis of the (TTAT)n repeat, as it is unstable in some families.

mothers and affected offspring with schizophrenia. We analysed 13 loci, which together span approximately 1.8 Mb of DNA and cover the class II and class III regions of the MHC. The order of these 13 loci is as follows: 6pter-AC004180-HSY14768-DJ201G24-HSA012008-HSMHC3W36A-HSMHC3A5-HS107715-HS172K2-HSDV19-HSEVMHC-HSO14-HS1033B10-HSICK721Q-6cen. The TDT showed that the HSMHC3A5 locus, lying near the junction of the class II and III regions, is significantly associated with schizophrenia (Table 1). The HSMHC3A5 locus contains the gene *NOTCH4*, spanning approximately 56.8 kb of DNA, inclusive of a putative promoter region and 30 exons⁶. We performed a further study using DNA marker haplotype systems consisting of three microsatellites and two SNPs within the HSMHC3A5 locus. Of the 5 markers, 3 are present in the 5' flanking region of *NOTCH4*, 1 in exon 1 and 1 in intron 17. Their order is as follows: 5'-(TAA)n-SNP1-SNP2-(CTG)n-(TTAT)n-3'. Of seven DNA marker haplotypes, the SNP2-(CTG)n haplotype showed the strongest association with schizophrenia ($P=0.0000078$), and the SNP1-SNP2-(CTG)n haplotype showed the second strongest association ($P=0.000011$; Table 2). To assess the effect of segregation distortion⁷, we also

analysed 48 unaffected offspring using these five markers. The TDT did not show significant distortion at these marker loci. Our results suggest that the susceptibility gene may be *NOTCH4* or a nearby locus.

The significant level of linkage disequilibrium (LD), extended over the (TAA)n repeat, but the (CT)n repeat in the testis-specific basic protein gene, adjacent to the 5' end of *NOTCH4* and present in the HS107715 locus, was not associated with schizophrenia. Moreover, four known genes (*PBX2*, *AGPAT1*, *AGER* and *TNXA*) and some ESTs encoding unknown proteins lie between *NOTCH4* and the HSMHC3W36A locus. Associations between these genes and schizophrenia cannot be ruled out, although the significant LD level did not extend over the (TTAT)n repeat because it was less informative than the (CTG)n and (TAA)n repeats.

The (CTG)n repeat in exon 1 of *NOTCH4* encodes leucine in the signal peptide domain. We have typed seven alleles using the (CTG)n repeat: (CTG)₅, (CTG)₆, (CTG)₉, (CTG)₁₀, (CTG)₁₁, (CTG)₁₂ and (CTG)₁₃. An excess of the (CTG)₁₀ allele was transmitted to affected offspring by their parents. Possibly, the (CTG)₁₀ allele itself may confer a susceptibility to schizophrenia. The SNP2, an A→G substitution in the promoter region, should also be considered as a candidate site.

Table 2 • Haplotype analysis for association of *NOTCH4* variation with schizophrenia

Haplotype	P value
(TAA)n-SNP1	0.0002
SNP1-SNP2	0.0162
SNP2-(CTG)n	0.0000078
(CTG)n-(TTAT)n	0.189
(TAA)n-SNP1-SNP2	0.000028
SNP1-SNP2-(CTG)n	0.000011
SNP2-(CTG)n-(TTAT)n	0.064

The (TAA)n repeat is 8.8 kb away from SNP1; SNP1 is present at base -1,725 of the 5' flanking region of *NOTCH4* and SNP2 at base -25; and the (CTG)n repeat is 40 bp away from SNP2 and 12.3 kb away from the (TTAT)n repeat.

Little is known about the functions of *NOTCH4* in humans, but animal studies demonstrate that the *NOTCH* family of proteins has a role in determining cell fate⁸. *NOTCH* was originally characterized as a *Drosophila melanogaster* neurogenic gene required for the correct segregation of epidermal cells from neuronal cell precursors during embryogenesis⁹. *In situ* hybridization in mice showed that although *Notch4* is highly expressed in endothelial cells, its transcripts are also detected in the developing nervous system¹⁰. These findings raise the possibility that *NOTCH4* may be involved in the development of neurogenic diseases.

Acknowledgements

We thank H. Hillman, R. Masters and I. Kilbourn for the collection of blood samples from family members; and A. Gliddon, P. Douglas, V. Hanson and L. Hughes for help.

Jun Wei & Gwynneth P. Hemmings

Institute of Biological Psychiatry, Schizophrenia Association of Great Britain, Bryn Hyfryd, The Crescent, Bangor, Gwynedd, UK.
Correspondence should be addressed to J.W. (e-mail: jwei@hgmp.mrc.ac.uk).

- Wang, S. et al. *Nature Genet.* 10, 41-46 (1995).
- Straub, R.E. et al. *Nature Genet.* 11, 235-236 (1995).
- Schwab, S.G. et al. *Nature Genet.* 11, 325-327 (1995).
- Schizophrenia Linkage Collaborative Group for Chromosomes 3, 6 and 8 *Am. J. Med. Genet.* 67, 580-694 (1996).
- Spielman, R.S. et al. *Am. J. Hum. Genet.* 52, 506-516 (1993).
- Li, L. et al. *Genomics* 51, 45-58 (1998).
- Spielman, R.S. & Ewens, W.J. *Am. J. Hum. Genet.* 59, 983-989 (1996).
- Song, W. et al. *Proc. Natl. Acad. Sci. USA* 96, 6959-6963 (1999).
- Fortini, M.E. & Artavanis-Tsakonas, S. *Cell* 75, 1245-1247 (1993).
- Uytenendaele, H. et al. *Development* 122, 2251-2259 (1996).
- Terwilliger, J.D. *Am. J. Hum. Genet.* 56, 777-787 (1995).